

**OPEN**

**Epidemiology Publish Ahead of Print**

**DOI: 10.1097/EDE.0000000000000858**

**Trends in Black and White Opioid Mortality in the United States, 1979–2015**

Monica J. Alexander<sup>1†</sup>; Mathew V. Kiang<sup>2†\*</sup>; Magali Barbieri<sup>1,3</sup>

<sup>1</sup> Department of Demography, University of California, Berkeley, Berkeley, California, USA

<sup>2</sup> Department of Social and Behavioral Sciences, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

<sup>3</sup> Institut national d'études démographiques, Paris, France

† These authors contributed equally to this work.

\* Corresponding author : mkiang@mail.harvard.edu, 617-384-8708, Dept Social and Behavioral Sciences, 401 Park Drive, 4<sup>th</sup> Floor, Boston, MA 02215

**Type of manuscript:** Original research article

**Conflicts of interest:** The authors have no conflicts of interest to declare.

**Sources of financial support:** MB was supported for this work by awards from the U.S.

National Institute on Aging (R01-AG011552 and R01-AG040245) as well as by contributions to the Human Mortality Database project from the Society of Actuaries in the United States and the United Kingdom Institute and Faculty of Actuaries. However, any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors alone and do not necessarily represent the official views of the National Institute on Aging and other funders.

MVK and MJA received no financial support for this work.

**Running head:** Trends in U.S. Opioid Mortality, 1979–2015

**Reproducibility:** All code to download and analyze the publicly-available data, as well as reproduce all tables and plots, is available at [https://github.com/mkiang/opioid\\_trends](https://github.com/mkiang/opioid_trends)

**Word count:** 3,107

Copyright © 2018 The Authors. Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ACCEPTED

## Abstract

### **Background:**

Recent research on the US opioid epidemic has focused on the white or total population and has largely been limited to data after 1999. However, understanding racial differences in long-term trends by opioid type may contribute to improving interventions.

### **Methods**

Using multiple cause of death data, we calculated age-standardized opioid mortality rates, by race and opioid type, for the US resident population from 1979 to 2015. We analyzed trends in mortality rates using joinpoint regression.

### **Results**

From 1979 to 2015, the long-term trends in opioid-related mortality for non-Hispanic blacks and non-Hispanic whites went through three successive waves. In the first wave, from 1979 to the mid-1990s, the epidemic affected both populations and was driven by heroin. In the second wave, from the mid-1990s to 2010, the increase in opioid mortality was driven by natural/semi-synthetic opioids (i.e., codeine, morphine, hydrocodone, or oxycodone) among whites, while there was no increase in mortality for blacks. In the current wave, increases in opioid mortality for both populations have been driven by heroin and synthetic opioids (i.e., fentanyl and its analogues). Heroin rates are currently increasing at 31% (95% CI: 27, 35) per year for whites and 34% (95% CI: 30, 40) for blacks. Concurrently, respective synthetic opioids are increasing at 79% (95% CI: 50, 112) and 107% (95% CI: -15, 404) annually.

## **Conclusion**

Since 1979, the nature of the opioid epidemic has shifted from heroin to prescription opioids for the white population to increasing of heroin/synthetic deaths for both black and white populations.

Key words: opioids; racial disparities; joinpoint analysis; substance abuse

ACCEPTED

## Introduction

Since 1979, the opioid mortality rate has increased 22-fold in the United States, resulting in over 33,000 deaths in 2015 alone, drawing national attention, and becoming an important part of the political agenda.<sup>1,2</sup> The continuous intensification of fatal overdoses is unique to opioids, as the rate of all non-opioid drug overdoses, though doubling from 1979 to 2006, has remained constant since then (eFigure 1; <http://links.lww.com/EDE/B356> and eTable 1; <http://links.lww.com/EDE/B356>).

In the 1990s and 2000s, the epidemic was driven by prescription opioids<sup>3,4</sup> as part of aggressive marketing tactics by the pharmaceutical industry and changing medical standards in diagnosis and treatment of chronic pain.<sup>5,6</sup> However, clinical behaviors (i.e., prescription patterns and pain management) differ markedly by race,<sup>7,8</sup> with much lower opioid prescription rates in the black population than in the white population.<sup>9,10</sup> In addition, the extent of racial bias in prescribing patterns has been found to vary by physician characteristics such as gender and race.<sup>11,12</sup> This differential clinician behavior is important as prescription opioid misuse is a substantial risk factor for heroin use.<sup>13</sup> Recent research has suggested that rates of heroin use and prescription opioid misuse in blacks and whites have begun to converge in recent years.<sup>14-16</sup> However, the convergence and divergence of opioid mortality rates by race over longer time periods have not been explored.

Recent research describes the opioid epidemic as having been initially driven by prescription opioids starting in the mid-1990s, followed by a continuing rise in the number of heroin deaths starting in the mid-2000s, concurrently fueled by illicitly manufactured synthetic opioids such as fentanyl starting around 2013.<sup>17-19</sup> However, these studies have largely been limited to opioid mortality in the total or white population after 1999, with only more recent

work focusing on the opioid epidemic in the black population.<sup>20</sup> We aim to supplement the current literature by describing black- and white-specific opioid mortality trends from 1979 to 2015. To date, this is the longest period of analysis of opioid mortality in the US, contributing an expanded perspective on trends over time and allowing for the examination of mortality by race and opioid type to understand the factors behind these trends.

## **Methods**

### **Multiple cause of death data**

We used a publicly available micro-level dataset with multiple causes of death derived from all the death certificates collected in the United States by the National Center for Health Statistics (NCHS) from 1979 to 2015.<sup>21</sup> For death certificates related to fatal overdoses, state laws require all contributing substances be identified, and where possible, a single agent to be specified as the underlying cause of death. In addition, up to 20 other contributory causes can be listed on the death certificate. The presence of a drug is not sufficient for the drug to be deemed the underlying cause; therefore, both the underlying and contributory causes must be considered when identifying deaths from drug overdose.<sup>4,22</sup>

Death certificates are coded by the NCHS to produce multiple cause of death data.<sup>21</sup> Over the period 1979–1998, deaths were classified according to the Ninth Revision of the International Classification of Diseases (ICD-9),<sup>23</sup> while the Tenth Revision (ICD-10)<sup>24</sup> has been implemented since 1999.

### **Defining opioid-related deaths**

We defined overdoses involving opioids as specified in the *Consensus recommendations for national and state poisoning surveillance* (eTable 2; <http://links.lww.com/EDE/B356>).<sup>25</sup> Specifically, for 1979–1998, an opioid overdose was defined as a death with at least one of the

following ICD-9 codes for either the underlying cause or for one of the contributory causes: E850.0 (heroin), E850.1 (methadone), E850.2 (other pharmaceutical opioid), or N965.0 (undetermined/unknown opioid).

For 1999–2015, all opioid overdoses were defined by a combination of specific codes for the underlying cause and at least one relevant contributory cause code. The ICD-10 codes for the underlying cause corresponding to overdoses are: X40–X44, X60–X64, X85 and Y10–Y14. In addition, for a death to be defined as opioid related, it has to include one or more of the following ICD-10 codes: opium (T40.0); heroin (T40.1); other natural and semi-synthetic (T40.2) such as codeine, morphine, hydrocodone, or oxycodone; methadone (T40.3); other synthetic (T40.4) such as tramadol, fentanyl, and fentanyl derivatives; or unspecified (T40.6).

### **Statistical analysis**

Using the multiple cause of death data, we tabulated deaths from opioid overdose for the total, black, and white resident populations by five-year age groups (i.e., 0-4, 5-9, ..., 85+). In addition, we tabulated these deaths by type of opioid. Due to the changes in the coding of opioid types from ICD-9 to ICD-10, we defined three temporally stable opioid categories as: (a) opioid deaths involving heroin, (b) opioid deaths involving methadone, and (c) opioid deaths involving an opioid other than (but not excluding) heroin or methadone. We refer to these categories as "heroin", "methadone", and "other" throughout the text; however, note that multiple opioids could be listed on death certificates coded with the ICD-10 and thus these categories are not mutually exclusive. The "other" category covers natural/semi-synthetic, synthetic, and unspecified opioid types. For the ICD-10 years (1999–2015), we additionally tabulated deaths by type of opioid using the more-specific ICD-10 categories defined above. To estimate the denominator of the rates, we used national race- and age-specific annual population estimates

from the US Census Bureau's Population Estimates Program. All rates are expressed as the number of deaths per 100,000 people and directly age-standardized to the US 2000 population. Standard errors for the rates were obtained using a Poisson approximation.<sup>26,27</sup> We calculated the opioid rate ratio as the rate for whites divided by the rate for blacks and estimated standard errors for the ratio using the delta approximation.<sup>28</sup>

We used joinpoint regression to examine trends in the age-standardized rates and rate ratio calculated as described above via the Joinpoint Regression Program version 4.5.0.1.<sup>29</sup> Using a segmented weighted least squares regression, the program iteratively adds joinpoints and finds the model with the optimal number of joinpoints, between zero and five, selecting the model that best fits the data. Model fit is evaluated using a permutation test.<sup>30</sup> To account for the ICD-9 to ICD-10 coding change, an indicator variable was used for years  $\geq 1999$  for models spanning the entire period (see eTables 3-8; <http://links.lww.com/EDE/B356> for full joinpoint model results). For the best model in each case, we report the annual percent change as well as the average annual percent change for the entire period under analysis. The average annual percentage change is defined as the weighted average of the annual percentage changes, with weights equal to the number of years in each time segment. Summary results are presented in Table 1 and Table 2. The code to reproduce tables and figures are available online (eText 1; <http://links.lww.com/EDE/B356>). This study was reviewed by the Harvard TH Chan School of Public Health institutional review board and did not require full review because it uses publicly available, retrospective, de-identified data.

## **Results**

From 1979 to 2015, the opioid mortality rate for whites increased from 0.44 to 12, corresponding to an average increase of 10% (95% CI: 9, 11) per year. Over the same period, the

rate for blacks increased from 0.62 to 6.6 for an average annual increase of 6% (95% CI: 5, 7). While the rate for whites increased steadily over the whole period, the rate for blacks remained stable in 1994–2011 (Table 1 and Figure 1). This differential growth is reflected in changes in the rate ratio (white to black mortality) through three successive waves. The rate ratio was approximately 0.71 from 1979 to 1993, then increased by 9% (95% CI: 8, 10) per year to over 2 by 2010 and declined thereafter by -6% (95% CI: -9, -3) annually to reach 1.9 in 2015 (Figure 1).

We examined trends in mortality by broad opioid category (heroin, methadone, and other) to identify the drivers of change in the rate ratio. Both blacks and whites experienced an increase in heroin and other opioid mortality over the full period (Table 1 and Figure 2). In 1979, the heroin mortality rate was 0.14 for whites and 0.33 blacks. By 2015, the heroin mortality rates had risen to 4.8 for whites and 3.1 for blacks, resulting in average annual increases of 12% (95% CI: 10, 14) for whites and 7% (95% CI: 6, 9) for blacks. The general pattern of increase in heroin mortality was similar for both populations: the rates first grew relatively slowly up to the 1990s, reaching a plateau in 1995 for blacks and 1998 for whites, before accelerating after 2005 for whites and 2010 for blacks. Between 2010 and 2015, the annual percent change in heroin mortality reached over 30% for both populations.

The mortality rate from other opioids increased from approximately 0.25 for each of the two populations to 8.2 for whites and 4.2 for blacks. In contrast to heroin mortality, the trends in other opioid mortality over the period differ markedly by race. For whites, the rate increased substantially in 1993–2010, while for blacks, it remained stable during this period. However, during 2013–2015, mortality due to other opioids increased substantially for both populations, at an annual rate of 18% (95% CI: 1, 39) for whites and 34% (95% CI: 2, 77) for blacks.

Opioid types can be further disaggregated during the period 1999–2015, due to the ICD-10 coding details (Table 2 and Figure 3). In addition to heroin and methadone, the ICD-10 also identifies opium (not analyzed due to low counts), natural/semi-synthetic opioids (e.g., oxycodone and hydrocodone), synthetic opioids other than methadone (e.g., fentanyl and tramadol), and unspecified opioids. The “other” opioid category defined above mainly consists of natural and semi-synthetic opioids and non-methadone synthetic opioids.

Trends in mortality due to natural/semi-synthetic opioids differ by race. By 2015, the natural opioid mortality rate for whites was more than double that for blacks, at 4.6 compared to 2.1. The rate for blacks increased steadily by 8% (95% CI: 8, 9) per year from 1999–2015. The white rate increased by 13% (95% CI: 12, 14) per year in 1999–2010, then leveled out in 2010–2015 at 1% (95% CI: -2, 3). The differential increase in natural opioid mortality by race explains the increase in the rate ratio from 1999–2010.

The synthetic opioid mortality rate was 0.30 for whites and 0.12 for blacks in 1999, and increased an average of 17% (95% CI: 13, 21) for whites and 15% (95% CI: 1, 32) for blacks. For whites, the synthetic mortality rate increased 79% per year (95% CI: 50, 112) from 2013–2015, largely due to fentanyl, often taken in combination with heroin.<sup>31</sup> While the confidence interval is wide and imprecise, the synthetic mortality rate for blacks for blacks was 107 (95% CI: -15, 404).

## **Discussion**

The opioid epidemic can be divided into three waves between 1979 and 2015. During the first wave, from 1979 to the mid-1990s, opioid mortality was higher for the black population but rates of increase were similar for both populations and largely driven by heroin. During the second wave, from the mid-1990s to 2010, the opioid epidemic expanded quickly within the

white population while opioid mortality remained stable in the black population. As a consequence, the racial gradient of risk reversed in 2000, and by 2010 the opioid mortality rate over 2 times higher for whites than for blacks. During this period, the opioid epidemic was driven largely by non-heroin and non-methadone opioids (i.e., prescription painkillers). Lastly, from about 2010 to 2015, the opioid mortality rate grew rapidly for both the black and white populations. This third wave has similarities to the first wave, as the mortality increase has been driven by heroin, and more recently synthetic opioids, in both populations. However, recent death rates due to heroin and synthetic opioids have reached unprecedented levels, increasing by an average of at least 30% per year since 2010 with further acceleration since 2013.

As reported in previous literature,<sup>4</sup> the acceleration in the mortality increase for whites since 1990 has been closely tied to an increase in the number of prescription drugs and the use of opioid pain medication, prompting several initiatives to curb the improper use of these painkillers. For example, in 2010, OxyContin was reformulated to be less easily abused. In addition, there have been other government-led initiatives such as Prescription Drug Monitoring Programs, Medicaid Lock-in programs and pain clinic laws, which sought to limit access and deter overuse of prescription opioids. These initiatives may explain the leveling of the natural/semi-synthetic opioid mortality rate that is observed since 2010.

The focus on restricting the supply of prescription opioids without concurrently reducing demand, however, appears to be associated with a shift from prescription painkillers to heroin and other illicit opioids. In recent years, the supply of both heroin and illegally manufactured fentanyl has increased in the United States.<sup>32,33</sup> The increased availability and affordability of illicit opioids has led to many users substituting heroin/fentanyl for prescription opioids. Previous research has demonstrated a clear pathway of addiction from prescription opioids to

heroin.<sup>34</sup> The Centers for Disease Control and Prevention reported that those addicted to opioid painkillers are 40 times more likely to be also addicted to heroin.<sup>35</sup>

However, the substitution of heroin for prescription opioids does not explain the recent rapid increase in the heroin and synthetic opioid mortality rates seen in the black population. Indeed, our findings show that the opioid mortality rate for blacks more than doubled in 5 years, even without a history of high prescription opioid mortality. Increased heroin mortality in more recent years is likely to be partly a consequence of the increased availability and affordability of heroin since 2010.<sup>36</sup> However, more recent increases in overdoses are likely to be driven by the increased potency of heroin, which is now often mixed with fentanyl.<sup>34</sup>

In addition to physician training and restrictions to the supply of prescription opioids, recent policies to curb the opioid epidemic have included comprehensive efforts such as harm reduction and early addiction treatment.<sup>37</sup> For example, access to naloxone, a drug designed to rapidly reverse the effects of opioid overdose, has been shown to reduce opioid mortality when expanded in settings where overdoses are more likely to occur.<sup>38,39</sup> Local interventions have expanded naloxone access by equipping law enforcement officers,<sup>40</sup> schools,<sup>41</sup> and libraries.<sup>42</sup> Similarly, there has been renewed interest in safe injection sites, where opioid users inject under the supervision of medical personnel,<sup>43</sup> which have been shown to reduce opioid overdose deaths<sup>44</sup> without an increase in drug trafficking or crime.<sup>45</sup> Effectively addressing the epidemic will continue to require a better understanding of social and economic determinants of opioid use and the impact of both supply-side and demand-side interventions.<sup>46,47</sup>

Our work contributes to the existing literature on understanding the opioid epidemic in three main ways. First, we presented an analysis of opioid mortality over an extended timeframe, analyzing an additional 20 years of data prior to 1999, when published studies usually begin.

This longer time series highlights similarities between historical conditions and the most recent wave of the epidemic. Second, we focused our analysis on differences in opioid mortality by race, whereas previous research tends to focus on trends in the total population, or in the white population only, thereby omitting some key racial differences in the epidemic over time. Finally, examining trends by type of opioid provides evidence of shifts in the epidemic from heroin to prescription drugs to heroin and fentanyl.

The current opioid epidemic in the United States is unprecedented compared to previous drug crises, in terms of the number of people affected and the rate of increase in overdose deaths. Our findings show that the nature of the epidemic has changed substantially over the past 35 years, shifting from relatively low heroin mortality, to a prescription drug problem in the white population, and more recently to a heroin/fentanyl epidemic affecting both the black and the white populations. In particular, since 2010, increases in mortality have been similar in both populations. The different patterns in the opioid epidemic by race suggest the need for targeted policy interventions to account for the distinctive pathways to addiction. A better understanding of racial differences and how they relate to the use of other drugs, place of residence, and socioeconomic status, is necessary to improve health interventions and rehabilitation programs across the country.

### **Limitations**

This study has several limitations. First, differences in cause-of-death reporting between blacks and whites could affect estimates of the rate ratio. Recent research has indicated that opioid mortality is underreported by about 24% nationally, and that the extent of underreporting varies by geography.<sup>48</sup> In particular, increases in heroin mortality were underreported in southern states such as Louisiana and Alabama, with large black populations, which may affect estimates

of heroin mortality among the black population, especially if this geographic effect is aggravated by further differential reporting by race. For example, if medical examiners are more likely to classify deaths of whites as due to an “unspecified drug” (ICD-10 code T50.9) rather than to opioids, the rate ratio estimate may be too conservative. Conversely, classifying black opioid deaths as an “unspecified drug” would make our rate ratio estimate too higher. The misclassification of race on death certificates could also affect the calculation of opioid mortality. However, while no validation studies have been carried out to examine the accuracy of coding of opioid deaths by race, an analysis of all causes of death showed that the effect on mortality of the misclassification of race on death certificates is small.<sup>49</sup>

Second, the jump joinpoint model assumes that the underlying trend remains the same during the ICD-9 to ICD-10 transition. While there are no validation studies on the effect of the ICD change on opioid deaths specifically, the comparability ratio of accidental injuries (under which accidental overdose — the overwhelming case of overdoses — is categorized) is estimated to be 1.03, which suggests that death classifications are reasonably comparable.<sup>50</sup> Unfortunately, the classification change occurred as the epidemic was accelerating and so quantifying this effect is difficult without further validation.

Last, it is important to note that fentanyl mortality rates may be underestimated, as a specific toxicology test must be requested by the coroner to determine the drug’s presence. Indeed, the observed increase in fentanyl deaths since 2013 is likely attributable, at least in part, to improved detection through post-mortem investigations due to increased awareness in the medical community.<sup>51</sup>

## **Conclusion**

Over the period 1979 to 2015, age-standardized opioid mortality rates increased for both the white and the black populations. While rates of increase were initially similar in the two populations and driven by heroin, the period 1993–2010 saw a rapid increase in opioid mortality for whites, largely due to prescription painkillers. Since 2010, opioid mortality has been increasing rapidly in both groups, largely driven by increases in heroin and synthetic opioids.

ACCEPTED

## References

1. White House. Obama Administration Announces Prescription Opioid and Heroin Epidemic Awareness Week. January 2016. <http://obamawhitehouse.archives.gov/the-press-office/2016/09/19/fact-sheet-obama-administration-announces-prescription-opioid-and-heroin>. Accessed July 4, 2017.
2. White House. President Donald J. Trump Signs an Executive Order Establishing the President's Commission on Combating Drug Addiction and the Opioid Crisis. January 2017. <https://www.whitehouse.gov/the-press-office/2017/03/30/president-donald-j-trump-signs-executive-order-establishing-presidents>. Accessed July 4, 2017.
3. Madras BK. The surge of opioid use, addiction, and overdoses responsibility and response of the US health care. *JAMA Psychiatry*. 2017;74(5):441-442. doi:10.1001/jamapsychiatry.2017.0163.
4. Paulozzi LJ, Budnitz DS, Xi Y. Increasing deaths from opioid analgesics in the United States. 2006;15(9):618-627. <http://doi.wiley.com/10.1002/pds.1276>.
5. Kolodny A, Courtwright DT, Hwang CS, et al. The Prescription Opioid and Heroin Crisis: A Public Health Approach to an Epidemic of Addiction. *Annu Rev Public Health*. 2015;36(1):559-574. doi:10.1146/annurev-publhealth-031914-122957.
6. Manchikanti L, Helm S, Fellows B, et al. Opioid epidemic in the United States. *Pain Physician*. 2012;15(3 Suppl):ES9-38. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=22786464&retmode=ref&cmd=prlinks>.
7. Pletcher MJ, Kertesz SG, Kohn MA, Gonzales R. Trends in opioid prescribing by race/ethnicity for patients seeking care in US emergency departments. *JAMA J Am Med*

- Assoc.* 2008;299(1):70-78.  
<http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2007.64>.
8. Netherland J, Hansen H. White opioids: Pharmaceutical race and the war on drugs that wasn't. *Biosocieties*. 2017;12(2):217-238. <https://link-springer-com.ezp-prod1.hul.harvard.edu/article/10.1057/biosoc.2015.46>.
  9. Anderson KO, Green CR, Payne R. Racial and Ethnic Disparities in Pain: Causes and Consequences of Unequal Care. *J Pain*. 2009;10(12):1187-1204. doi:10.1016/j.jpain.2009.10.002.
  10. Singhal A, Tien Y-Y, Hsia RY. Racial-Ethnic Disparities in Opioid Prescriptions at Emergency Department Visits for Conditions Commonly Associated with Prescription Drug Abuse. Seedat S, ed. *PLoS One*. 2016;11(8):e0159224. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0159224>.
  11. Burgess DJ, Phelan S, Workman M, et al. The effect of cognitive load and patient race on physicians' decisions to prescribe opioids for chronic low back pain: A randomized trial. *Pain Med (United States)*. 2014;15(6):965-974. doi:10.1111/pme.12378.
  12. Chapman EN, Kaatz A, Carnes M. Physicians and implicit bias: How doctors may unwittingly perpetuate health care disparities. *J Gen Intern Med*. 2013;28(11):1504-1510. doi:10.1007/s11606-013-2441-1.
  13. Compton WM, Jones CM, Baldwin GT. Relationship between Nonmedical Prescription-Opioid Use and Heroin Use. *N Engl J Med*. 2016;374(2):154-163. doi:10.1056/NEJMra1508490.
  14. Pouget ER, Fong C, Rosenblum A. Racial/Ethnic Differences in Prevalence Trends for Heroin use and Non-Medical use of Prescription Opioids Among Entrants to Opioid

- Treatment Programs, 2005–2016. *Subst Use Misuse*. 2017;1-11.  
doi:10.1080/10826084.2017.1334070.
15. Martins SS, Santaella-Tenorio J, Marshall BDL, Maldonado A, Cerd?? M. Racial/ethnic differences in trends in heroin use and heroin-related risk behaviors among nonmedical prescription opioid users. *Drug Alcohol Depend*. 2015;151:278-283.  
doi:10.1016/j.drugalcdep.2015.03.020.
  16. Metz VE, Sullivan MA, Jones JD, et al. Racial Differences in HIV and HCV Risk Behaviors, Transmission, and Prevention Knowledge among Non-Treatment-Seeking Individuals with Opioid Use Disorder. *J Psychoactive Drugs*. 2017;49(1):59-68.  
doi:10.1080/02791072.2016.1259518.
  17. Frenk SM, Porter KS, Paulozzi LJ. Prescription opioid analgesic use among adults: United States, 1999-2012. *NCHS Data Brief*. 2015;(189):1-8.  
<http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=25714043&retmode=ref&cmd=prlinks>.
  18. Dart RC, Surratt HL, Cicero TJ, et al. Trends in Opioid Analgesic Abuse and Mortality in the United States. January 2015. <http://www.nejm.org.ezp-prod1.hul.harvard.edu/doi/full/10.1056/NEJMsa1406143>.
  19. Rudd RA, Aleshire N, Zibbell JE, Gladden RM. Increases in Drug and Opioid Overdose Deaths--United States, 2000-2014. *MMWR Morb Mortal Wkly Rep*. 2016;64(50-51):1378-1382. <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6450a3.htm>.
  20. Bechteler SS, Kane-Willis K. Whitewashed : The African American Opioid Epidemic. 2017;(November).  
<https://www.thechicagourbanleague.org/cms/lib/IL07000264/Centricity/Domain/1/White>

washed AA Opioid Crisis 11-15-17\_EMBARGOED\_FINAL.pdf.

21. National Center for Health Statistics. *Data Access — Vital Statistics Online (Mortality Multiple Cause Files)*. [http://www.cdc.gov/nchs/data\\_access/vitalstatsonline.htm](http://www.cdc.gov/nchs/data_access/vitalstatsonline.htm), accessed June 27 2017.; 2017.
22. Stephens BG, Jentzen JM, Karch S, Wetli C V, Mash DC. National Association of Medical Examiners Position Paper on the Certification of Cocaine-Related Deaths. 2004;25(1):11-13.  
<http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00000433-200403000-00002>.
23. WHO. *International Classification of Diseases; Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death: Ninth Revision (ICD-9)*. Geneva, 1977.; 1977.
24. WHO. *International Classification of Diseases; Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death: Tenth Revision (ICD-10)*. Geneva, Switzerland; 1997. <http://apps.who.int/classifications/icd10/browse/2016/en>.
25. ISW7. *Consensus Recommendations for National and State Poisoning Surveillance*. Atlanta, Georgia: The Safe States Alliance; 2012.
26. Curtin LR, Klein RJ. Direct standardization (age-adjusted death rates). *Health People 2000 Stat notes*. 1995;(6):1-10.  
<http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=11762384&retmode=ref&cmd=prlinks>.
27. Keyfitz N. Sampling variance of standardized mortality rates. 1966;38(3):309-317.
28. Flanders WD. Approximate variance formulas for standardized rate ratios. *J Chronic Dis*.

- 1984;37(6):449-453. doi:10.1016/0021-9681(84)90028-6.
29. Statistical Methodology and Applications Branch Surveillance Research Program. Joinpoint Regression Program. 2017.
  30. Kim HJ, Fay MP, Feuer EJ, Midthune DN. Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med*. 2000;19(3):335-351. doi:10.1002/(SICI)1097-0258(20000215)19:3<335::AID-SIM336>3.0.CO;2-Z.
  31. O'Donnel KJ, Gladden RM, Seth P. Trends in Deaths Involving Heroin and Synthetic Opioids Excluding Methadone, and Law Enforcement Drug Product Reports, by Census Region — United States, 2006–2015. 2017;66:879-903. doi:10.15585/mmwr.mm6634a2.
  32. Office of National Drug Control Policy. *National Drug Control Strategy: Data Supplement*. [http://obamawhitehouse.archives.gov/sites/default/files/ondcp/policy-and-research/2015\\_data\\_supplement\\_final.pdf](http://obamawhitehouse.archives.gov/sites/default/files/ondcp/policy-and-research/2015_data_supplement_final.pdf), accessed August 13 2017.; 2015.
  33. Centers for Disease Control and Prevention. Increases in Fentanyl Drug Confiscations and Fentanyl-related Overdose Fatalities. 2015.
  34. Cicero TJ, Ellis MS, Kasper ZA. Increased use of heroin as an initiating opioid of abuse. *Addict Behav*. 2017;74:63-66. doi:10.1016/j.addbeh.2017.05.030.
  35. Centers for Disease Control and Prevention. *Today's Heroin Epidemic: Vital Signs*. <http://www.cdc.gov/vitalsigns/heroin/index.html>; 2015.
  36. Center for Behavioral Health Statistics and Quality. Behavioral health trends in the United States: Results from the 2014 National Survey on Drug Use and Health. *HHS Publication No SMA 15-4927, NSDUH Ser H-50*. 2015:64. <http://www.samhsa.gov/data/sites/default/files/NSDUH-FRR1-2014/NSDUH-FRR1-2014.pdf>.

37. Kolodny A, Frieden TR. Ten steps the federal government should take now to reverse the opioid addiction epidemic. *JAMA*. October 2017. doi:10.1001/jama.2017.14567.
38. Walley AY, Xuan Z, Hackman HH, et al. Opioid overdose rates and implementation of overdose education and nasal naloxone distribution in Massachusetts: Interrupted time series analysis. *BMJ*. 2013;346(7894). doi:10.1136/bmj.f174.
39. Coffin PO, Sullivan SD. Cost-effectiveness of distributing naloxone to heroin users for lay overdose reversal. *Ann Intern Med*. 2013;158(1):1-9. doi:10.1016/j.jemermed.2013.03.002.
40. Davis CS, Carr D, Southwell JK, Beletsky L. Engaging Law Enforcement in Overdose Reversal Initiatives: Authorization and Liability for Naloxone Administration. *Am J Public Health*. 2015;105(8):1530-1537. <http://dx.doi.org/10.2105/AJPH.2015.302638>.
41. Montgomery County Public Schools. *Policy and Regulations: IGN, Preventing Alcohol, Tobacco, and Other Drug Abuse in Montgomery County Public Schools*. <http://www.montgomeryschoolsmd.org/departments/policy/detail.aspx?recID=208&policyID=IGN&sectionID=9>, accessed August 13 2017.; 2017.
42. Ford A. *Saving Lives in the Stacks: How Libraries Are Handling the Opioid Crisis*. <https://americanlibrariesmagazine.org/2017/06/21/saving-lives-in-the-stacks/>, accessed August 13 2017.; 2017.
43. Miller CL, Kerr T, Frankish JC, et al. Binge drug use independently predicts HIV seroconversion among injection drug users: implications for public health strategies. *Subst Use Misuse*. 2006;41(2):199-210. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=16393742&retmode=ref&cmd=prlinks>.

44. Marshall BD, Milloy MJ, Wood E, Montaner JS, Kerr T. Reduction in overdose mortality after the opening of North America's first medically supervised safer injecting facility: A retrospective population-based study. *Lancet*. 2011;377(9775):1429-1437. doi:10.1016/S0140-6736(10)62353-7.
45. Potier C, Lapr evote V, Dubois-Arber F, Cottencin O, Rolland B. Supervised injection services: What has been demonstrated? A systematic literature review. *Drug Alcohol Depend*. 2014;145:48-68. doi:10.1016/j.drugalcdep.2014.10.012.
46. Dasgupta N, Beletsky L, Ciccarone D. Opioid Crisis: No Easy Fix to Its Social and Economic Determinants. 2018;108(2):182-186. doi:10.2105/AJPH.2017.304187.
47. Diez Roux A V. Despair as a cause of death: More complex than it first appears. *Am J Public Health*. 2017;107(10):1566-1567. doi:10.2105/AJPH.2017.304041.
48. Ruhm CJ. Geographic Variation in Opioid and Heroin Involved Drug Poisoning Mortality Rates. *American Journal of Preventive Medicine*. 2017.
49. Arias E, Heron M, National Center for Health S, Hakes J, Bureau USC. The Validity of Race and Hispanic-origin Reporting on Death Certificates in the United States: An Update. *Vital Heal Stat 2*. 2016;(172):1-21. <https://www.ncbi.nlm.nih.gov/pubmed/28436642>.
50. Anderson RN, Mini no a M, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: preliminary estimates. *Natl Vital Stat Rep*. 2001;49(2):1-32.
51. Slavova S, Costich JF, Bunn TL, et al. Heroin and fentanyl overdoses in Kentucky: Epidemiology and surveillance. *Int J Drug Policy*. 2017;46:120-129. doi:10.1016/j.drugpo.2017.05.051.

## Figures

Figure 1. A) Age-standardized opioid mortality rate for the white (red), and black (blue) US resident populations, 1979–2015. B) Rate ratio (white / black) of opioid mortality rates. Dots are estimated rate. Vertical bars represent 95% confidence interval. Solid lines are joinpoint model fits.

Figure 2. Age-standardized opioid mortality rates by general type of opioid: heroin (green), methadone (orange) or other opioid (violet) for white (top) and black (bottom) US resident populations, 1979–2015. Vertical bars are 95% confidence intervals. Solid lines are joinpoint model fits.

Figure 3. Opioid mortality rates for ICD-10 years (1999–2015) by type for white (top) and black (bottom) US resident populations. Opioid types include heroin (green), methadone (orange), natural/semi-natural opioids (purple), synthetic opioids (pink), and unspecified opioids (light green). Vertical bars are 95% confidence intervals. Solid lines are joinpoint model fits.



**Table 2.** Joinpoint Regression Results by Race and ICD-10 Opioid Type, 1999–2015.

	Observed value <sup>a</sup>				Average APC		Trend 1			Trend 2			Trend 3		
	1999		2015		AAPC	95% CI	Years	APC	95% CI	Years	APC	95% CI	Years	APC	95% CI
Heroin															
White	0.74	0.71, 0.78	4.8	4.7, 4.9	13	10, 16	1999-2005	1	-4, 5	2005-2010	12	5, 21	2010-2015	31	27, 35
Black	0.83	0.73, 0.93	3.1	2.9, 3.3	9	7, 11	1999-2010	-1	-3, 2	2010-2015	34	30, 40			
Methadone															
White	0.31	0.29, 0.34	1.2	1.2, 1.2	9	7, 12	1999-2003	43	34, 52	2003-2006	21	8, 36	2006-2015	-6	-7, -5
Black	0.21	0.16, 0.26	0.58	0.51, 0.65	6	4, 8	1999-2006	22	17, 27	2006-2015	-5	-6, -3			
Natural															
White	1.1	1.1, 1.2	4.6	4.5, 4.7	9	8, 10	1999-2010	13	12, 14	2010-2015	1	-2, 3			
Black	0.62	0.54, 0.71	2.1	2.0, 2.3	8	8, 9	1999-2015	8	8, 9						
Synthetic															
White	0.30	0.28, 0.32	3.6	3.5, 3.7	17	13, 21	1999-2006	18	11, 24	2006-2013	3	-2, 8	2013-2015	79	50, 112
Black	0.12	0.078, 0.15	2.1	2.0, 2.2	15	1, 32	1999-2013	6	-6, 19	2013-2015	107	-15, 404			
Unspecified															
White	0.96	0.92, 1.0	1.0	1.0, 1.0	0	-1, 0	1999-2015	0	-1, 0						
Black	2.1	1.9, 2.2	0.59	0.52, 0.67	-8	-9, -7	1999-2015	-8	-9, -7						

<sup>a</sup>All observed values are expressed as rates per 100,000; APC, annual percent change; AAPC, average annual percent change; CI, confidence interval





